THE 2010–2011 KENNESAW STATE UNIVERSITY
HIGH SCHOOL MATHEMAT ICS COMPETITION

PART I – MULTIPLE CHOICE

For each of the following 25 questions, carefully blacken the appropriate box on the answer sheet with a #2 pencil. Do not fold, bend, or write stray marks on either side of the answer sheet. Each correct answer is worth 6 points. Two points are given if no box, or more than one box, is marked. Zero points are given for an incorrect answer. Note that wild guessing is apt to lower your score. When time is over, give your answer sheet to your proctor. You may keep your copy of the questions.

NO CALCULATORS

90 MINUTES

1. Although Elvis Presley had more top ten hits than the Beatles (Elvis had 38, the Beatles had 33), the Beatles had two more of these top ten hits reach number one than Elvis had. If between them there were a total of 33 songs that reached the top ten but not number one, how many number one hits did the Beatles have?

   (A) 14 (B) 16 (C) 18 (D) 20 (E) 22

2. A date is called weird if the number of its month and the number of its day are relatively prime (have greatest common factor 1). For example, April 21 is weird because 4 and 21 are relatively prime. What is the smallest number of weird days in any month?

   (A) 15 (B) 14 (C) 13

3. In magic squares, each row, column, and diagonal must add up to what number?

2	3	
		4
		1

   (A) 4.25 (B) 4.5 (C) 5 (D) 5.25 (E) 6.5

4. If L has equation $ax + by = c$, M is its reflection across the x-axis, and N is its reflection across the y-axis, which of the following must be true about M and N for all nonzero choices of a, b, and c?

   (A) the x-intercepts are equal
   (B) the y-intercepts are equal
   (C) the slopes are equal
   (D). the slopes are reciprocals
   (E) the slopes are negative reciprocals

5. The counting numbers are written in the pattern at the right. What is the middle number of the 40th row?

```
          1
       2  3  4
    5  6  7  8  9
10 11 12 13 14 15 16
```

   (A) 1561 (B) 1641 (C) 1559 (D) 1639 (E) 1483

12. Twelve lattice points are arranged along the edges of a 3×3 square as shown. How many triangles have all three of their vertices among these points? One such triangle is shown.



(A) 48 (B) 64 (C) 204 (D) 220 (E) 256

13. The year 2010 has three consecutive prime number factors, 2, 3, and 5. How many years in the third millenium, i.e. between the years 2001 and 2999, have

20. Suppose f(x) = ax + b, g(x) = bx + a (a, b integers). If f(1) = 8 and f(g(50)) - g(f(50)) = 28, find the product of a and b.

(A) 5 (B) 12 (C) 48 (D) 182 (E) 210

21. One of the roots of the equation $x^3 - 14x^2 + 26x + c = 0$ is the quotient of the other two roots. Compute the product of all possible values of c.

(A) 456 (B) 529 (C) 576 (D) 625 (E) 676

22. In the rectangular solid, the areas of the following rectangles

SOLUTIONS – KSU MATHEMATICS COMPETITION – 2010

1. D Let B = number of #1 hits for the Beatles, Let E = number of #1 hits for Elvis. Then B = E + 2 and 33 = B + 38 – E = 33. Solving gives E = 18 and B = 20.

2. E Clearly, the number of the month in question must have the largest possible number of prime factors in order to minimize the number days with which it has a common factor other than 1. These are months 6, 10, and 12 (June, October, and December). Each has two prime factors, but October and December have 31 days, and 31 is a prime number. So let's consider June first. There are 10 days in June whose greatest common factor with 6 is 1 (1, 5, 7, 11, 13, 17, 19, 23, 25, 29) [There are 13 days in October whose greatest common factor with 10 is 1 (1, 3, 7, 9, 11, 13, 17, 19, 21, 23, 27, 29, and 31). There are 11 days in December whose greatest common factor with 12 is 1 (1, 5, 7, 11, 13, 17, 19, 23, 25, 29, and 31).] Therefore, the correct month is June and the number of "weird" days is 10.

3. B There are several ways to proceed. Here is one approach.

2	3	X
	B	4
A	C	1

The desired sum is X + 5.
Since 2+B+1 = X+5, B = X+2.
Since A+B+X = A + X + 2 + X = X + 5, A = 3 – X
Since 3 + B + C = 3 + X + 2 + C = X + 5, C = 0
Thus, the magic square becomes:

2	3	X
	X+2	4
3-X	0	1

Summing the bottom row, 3 – X + 0 + 1 = X + 5
which yields X = 5. Therefore, each row, column, and main diagonal has a sum of 10.

4. C Even a quick rough sketch like the one shown demonstrates that the reflection images are parallel and thus have the same slope.
Proof: Reflection preserves angle measurement. From the diagram, D + E = 90, therefore 2D + 2E = 180 so that the same-side interior angles of M and N are supplementary, proving the lines parallel.



5. A Note that the number on the right of each row is a perfect square, so that the 40th row ends in $40^2 = 1600$. Also note that the first number in the 40th row is $39^2 + 1 = 1522$. Since each row has an odd number of numbers, the middle number of the 40th row is the average of the first and last or 1561.

6. C When a = 0, $x = \frac{c}{b}$. We know $2(2^2) + 2b + c = 0 \Rightarrow 2b + c = -8$. We also know $1 + b + c = 0 \Rightarrow b + c = -1$. Solving the two equations together, b = -7 and c = 6. Therefore, when a = 0, $x = \frac{6}{7}$.

13. A Let us first list the sets of three consecutive primes whose product is

17. E Let S denote the given sum. Since $\cos(90° - x) = \sin(90° - x)$ and $\sin^2(x) = 1 - \cos^2(x)$, we get $S = \cos^2(1°) + \cos^2(2°) + \cos^2(3°) + \ldots + \cos^2(90°)$

$= \sin^2(90° - 1°) + \sin^2(90° - 2°) + \sin^2(90° - 3°) + \ldots + \sin^2(90° - 89°)$

$= \sin^2(89°) + \sin^2(88°) + \sin^2(87°) + \ldots + \sin^2(1°)$

$= (1 - \cos^2(89°)) + (1 - \cos^2(88°)) + (1 - \cos^2(87°)) + \ldots + (1 - \cos^2(1°)) = 89 - S$

Thus $2S = 89$, which implies that $S = 44.5$.

18. D Represent the first six terms of the geometric sequence as $a, ar, ar^2, ar^3, ar^4, ar^5$.
Then $a + ar + ar^2 + ar^3 + ar^4 + ar^5 = 91$ and $a + ar = 7$.
Grouping the left side of the first equation in pairs, and factoring:

$a + ar + ar^2 + ar^3 + ar^4 + ar^5 = (a + ar)(1 + r^2 + r^4)$.

Thus, $7(1 + $ [illegible] ^{2}

22. D Solving the equations $xy = 300$, $xz = 240$ and $yz = 320$ simultaneously gives the dimensions of the rectangular solid: $x = 15$, $y = 20$, and $z = 16$.

Using the Pythagorean Theorem on right triangles HEA, HGC, and ABC we find $HA = \sqrt{481}$, $HC = \sqrt{656}$, and $AC = 25$.

Representing HK as p, AK as a and KC as $(25 - a)$, and using the Pythagorean Theorem on right triangles HAK and HCK,

$$a^2 + p^2 = 481 \quad \text{and} \quad (25 - a)^2 + p^2 = 656$$

Solving these two equations together gives $a = 9$ and $p = 20$.
Noting that triangle GCK is an isosceles right triangle with legs of 16, $GK = 16\sqrt{2}$.

Since $HG = HK = 20$, $\triangle GHK$ is isosceles. Its area can be found by drawing an altitude to GK (length $\sqrt{272}$) and using area = ½bh.
The area of $\triangle GHK$ is $32\sqrt{34}$.

23. E Let g be the number of green beads in a box. Then the total number of beads in that box is $g + 6$. The number of ways of drawing two beads together from the box is ${}_{(g+6)}C_2 = \dfrac{(g+6)(g+5)}{2}$. The number of ways of drawing together 1 green and 1 blue bead is $6g$. Thus, the probability of drawing 1 of each color is

$$\frac{6g}{\dfrac{(g+6)(g+5)}{2}} = \frac{12g}{(g+6)(g+5)} =$$

25. B The following lengths can be determined fairly easily:
EC = 1, CD = 3, AD = 2, BF = 1, EB = 4, AF = 5.

Also using right triangle ABC, sin