


**THE 2013-2014 KENNESAW STATE UNIVERSITY
HIGH SCHOOL MATHEMATICS COMPETITION
PART II**

In addition to scoring student responses based on whether a solution is correct and complete, consideration will be given to elegance, simplicity, originality, and clarity of presentation.

Calculators are NOT permitted.

- !
1. A and B both represent nonzero digits (not necessarily distinct). If the base ten numeral $\underline{A}\underline{B}$ divides, without remainder, the base ten numeral $\underline{A}0\underline{B}$ (whose middle digit is zero), find, with proof, all possible values of $\underline{A}\underline{B}$.
- !


2. A and B are points on the positive x and positive y axes respectively and C is the point with coordinates $(3, 4)$. Prove that the perimeter of triangle ABC is greater than 10.



3. One solution for the equation $a^2 + b^2 + c^2 + 2 = abc$ is $a = 3, b = 3$ and $c = 4$.
- Find a solution (a, b, c) where $a, b,$ and c are integers all larger than 10.
 - Prove that there are infinitely many solutions (a, b, c) where $a, b,$ and c are positive integers.

4. Consider the equation $\sqrt{x} = \sqrt{a} + \sqrt{b}$, where x is a positive integer.
- Prove that the equation has a solution (a, b) where a and b are both positive integers, if and only if x has a factor which is a perfect square greater than 1.
 - If $x \neq 1,000$, compute, with proof, the number of values of x for which the equation has at least one solution (a, b) where a and b are both positive integers.

5. In right triangle ABC , $AC = 6, BC = 8$ and $AB = 10$. PA and PB bisect angles A and B respectively. Compute, with proof, the ratio $\frac{PA}{PB}$.



1. Of course, this problem can be done by trial and error (there are only 81 possibilities), but we present a more elegant solution.

Suppose $\frac{AB}{AB}$

3. Suppose we begin with two positive integers a and b , and we try to find a third integer x such that $a^2 + b^2 + x^2 + 2 = abx$. Then the problem can be thought of as finding an integer solution (if one exists) for the quadratic equation $(ab)x + (a^2 + b^2 + 2) = 0$.

If there is some integer solution $x = c$, then there must exist a real number d such that

$$x^2 - (ab)x + (a^2 + b^2 + 2) = (x - c)(x - d) = x^2 - (c+d)x + cd$$

Comparing the coefficients on the left and right sides of this last equation, we know that $ab = c + d$, so that $d = ab - c$ is also an integer. Therefore, given any three integers a , b , and c such that $a^2 + b^2 + c^2 + 2 = abc$, we can replace c with $ab - c$ to obtain another solution.

We know that $(4, 3, 3)$ is a solution. So we can replace one of the 3's with $4 \cdot 3 - 3 = 9$ to get the solution $(4, 3, 9)$. Since a , b , and c are interchangeable, we can obtain other solutions by repeatedly replacing the smallest number (which we will call $ab - c$). Hence, listing the numbers in decreasing order at each step, we obtain the following solutions:

$$(4, 3, 3) \rightarrow (9, 4, 3) \rightarrow (33, 9, 4) \rightarrow (293, 33, 9) \rightarrow (9660, 293, 33).$$

Since this process can be repeated indefinitely, there are infinitely many positive integer solutions (a, b, c) to the given equation.

4. (i) Given $\sqrt{x} = \sqrt{a} + \sqrt{b}$.

Suppose $x = ky^2$, with k and y positive integers, and $k > 1$. We must prove that there exists at least one pair of positive integers (a, b) that satisfies the equation.

We have $\sqrt{x} = \sqrt{k^2 y} = k\sqrt{y}$. Since $k > 1$, then $k - 1 > 0$. Therefore,

$$\sqrt{x} = k\sqrt{y} = (k-1)\sqrt{y} + \sqrt{y} = \sqrt{(k-1)^2 y} + \sqrt{y}.$$

Since both $(k-1)^2 y$ and y are both positive integers, setting $a = (k-1)^2 y$ and $b = y$ gives the desired result.

5. Method 1:

We will refer to $\angle CAB$ as $\angle A$ and $\angle CBA$ as $\angle B$.
So that $m\angle A + m\angle B = 90^\circ$.

Then $m\angle P = 180^\circ - (m\angle A + m\angle B) = 135^\circ$.
So that, $m\angle PAB + m\angle PBA = 45^\circ$. Represent the measures of these two angles with θ and $45^\circ - \theta$.

Using the Law of Sines on $\triangle APB$

$$\frac{PA}{\sin(45^\circ - \theta)} = \frac{\sin 45^\circ \cos \theta + \cos 45^\circ \sin \theta}{\sin \theta} = \sin 45^\circ \cot \theta + \cos 45^\circ.$$

Now $\cot \theta = \cot(\angle A) = \frac{1 + \cos A}{\sin A}$ (using the appropriate half-angle formula)


But in $\triangle ABC$, $\cos A = \frac{6}{10}$ and $\sin A = \frac{8}{10}$, making $\cot \theta = \frac{1 + \frac{6}{10}}{\frac{8}{10}} = 2$.

Finally, $\frac{PA}{PB} = (\sin 45^\circ)(2) + \cos 45^\circ = \frac{\sqrt{2}}{2}(2) + \frac{\sqrt{2}}{2} = \frac{3\sqrt{2}}{2}$.

Method 2:

Note that since point P is the intersection of the angle bisectors of $\triangle ABC$, P is the incenter (the center of the inscribed circle).

Noting that the tangent segments to a circle from an external point are congruent, represent the lengths of the segments in the diagram as shown.



Then $6 - x + 8 - x = 10$ and $x = 2$.

Therefore, right $\triangle ARP$ has side lengths 2, 4, and $2\sqrt{5}$, and right $\triangle BMP$ has side lengths 2, 6, and $2\sqrt{10}$.

Therefore, $\frac{PA}{PB} = \frac{2\sqrt{5}}{2\sqrt{10}} = \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2}$.